



2. Advanced pharmacological skills: Venipuncture/vascular access	
i. Vaccine administration	<b>P</b> <b>I*</b> <b>A*</b> <b>C*</b> <b>P</b> EMT-Is, AEMTs and CTs may only administer vaccinations during designated events such as mass vaccination clinics or in the event of a declared public health emergency and then only after approved training.
j. Paralytic administration	<b>P*</b> Administration of paralytics for the purpose of RSI is not permitted unless the EMS agency has met the RSI requirements promulgated by OEMS and has received approval for RSI use from OEMS. Paramedics are authorized to use paralytics to maintain the paralysis of already intubated patients, if approved by the local EMS Medical Director.
k. Administration other physician approved medications	<b>P*</b> <b>C*</b> <b>P*</b> CTs are only authorized to give the following: anti-arrhythmics, vagolytic agents, chronotropic agents, alkalinizing agents, analgesic agents and vasopressor agents. Paramedics are authorized to give any medication via enteral or parenteral routes, if approved by the local EMS Medical Director.
l. Maintain an infusion of blood or blood products	<b>P</b>

Cardiac /Medical Skills		Levels		Interpretive Guidelines	
1. Fundamental cardiac skills					
a. Manual external CPR	<b>E</b> <b>I</b> <b>A</b> <b>C</b> <b>P</b>	<b>E</b> <b>I</b> <b>A</b> <b>C</b> <b>P</b>	<b>E</b> <b>I</b> <b>A</b> <b>C</b> <b>P</b>		
b. Use of an automated external defibrillator	<b>E</b> <b>I</b> <b>A</b> <b>C</b> <b>P</b>	<b>E</b> <b>I</b> <b>A</b> <b>C</b> <b>P</b>	<b>E</b> <b>I</b> <b>A</b> <b>C</b> <b>P</b>		
2. Advanced cardiac skills					
a. Use mechanical CPR assist devices	<b>E</b> <b>I</b> <b>A</b> <b>C</b> <b>P</b>	<b>E</b> <b>I</b> <b>A</b> <b>C</b> <b>P</b>	<b>E</b> <b>I</b> <b>A</b> <b>C</b> <b>P</b>		
b. ECG monitoring and interpretation	<b>E*</b> <b>I*</b> <b>A*</b> <b>C</b> <b>P</b>	<b>E*</b> <b>I*</b> <b>A*</b> <b>C</b> <b>P</b>	<b>E*</b> <b>I*</b> <b>A*</b> <b>C</b> <b>P</b>		Includes 12-lead ECGs. EMTs, EMT-Is, and AEMTs may only obtain and transmit a 12-lead ECG for suspected STEMI patients, if approved and trained by the local EMS Medical Director. ECG interpretation is limited to CTs and Paramedics.

EMT
E
EMT-I
I
AEMT
A
CT
C
PMDC
P