


Criminal Justice Georgia Rate



LC-MS/MS Confirmations/Determinations - Urine Continued

Other LC-MS/MS Tests	Other LC-MS/MS Determinations						
Cyclobenzaprine	each	\$8.80	D/L Methamphetamine	each	\$39.00		
Diphenhydramine	each	\$8.80	Designer Drugs (a-PVP, Butylone, Ethylone,	panel	\$25.00		
Duloxetine (Cymbalta)	each	\$8.80	MDPV, Mephedrone, Methylone)		,		
Gabapentin	each	\$8.80	GHB	each	\$34.00		
Methylphenidate (Ritalin)	each	\$8.80	Ketamine	each	\$11.55		
Naltrexone	each	\$8.80	Kratom (Mitragynine)	each	\$44.00		
Pregabalin	each	\$8.80	LSD	each	\$55.00		
Sertraline (Zoloft)	each	\$8.80	Nicotine/Cotinine	each	\$6.60		
Tapentadol	each	\$8.80					
Trazadone (as mCPP)	each	\$8.80	LC-MS/MS	each	\$8.80		
Zolpidem (Ambien)	each	\$8.80	Determination not otherwise listed				

QTOF Analysis - Urine Qualitative (Positive/Negative) Results

Comprehensive Panel (104 analytes) \$45.50
Designer Stimulants Panel (20 Drug) \$28.00
Synthetic Cannabinoids Panel (19 analogs) \$25.00

Oral Fluid Tests - LC-MS/MS only - Select analytes to customize a panel

	Base Price	e - Collection Tube	each \$4	.00		
Amphetamine Barbiturate Benzodiazepine Buprenorphine Methadone Opiates Oxycodone	each each each each each each	\$6.60 \$6.60 \$6.60 \$6.60 \$6.60 \$6.60 \$6.60	Illicit Drugs Methamph Cocaine Ecstasy Phencyclic THC	hetamine - Includ	ded in Amphetamine panel each each each each	\$6.60 \$6.60 \$6.60 \$6.60
Synthetic Opiates			Other			
Fentanyl Meperidine Propoxyphene Tramadol	each each each each	\$6.60 \$6.60 \$6.60 \$6.60	Carisoprod Tapentado		each each	\$6.60 \$6.60

Pharmacogenetic (PGx) Testing Quantity Price
--

Precision Medication Management - Psychiactric

each

\$300.00

Up to 95% of drug variability between individuals is due to genetic factors. PGx testing can help shorten response time by limiting trial-and-error prescribing. In addition, metabolically inappropriate drugs and their side effects can be avoided, while drugs more likely to be effective are identified as clear, first-choice options.

^{*} For best results, provide all medications patient is taking or could be prescribed.